

THE PECULIAR BALMER DECREMENT OF SN 2009IP: CONSTRAINTS ON CIRCUMSTELLAR GEOMETRY

EMILY M. LEVESQUE¹, GUY S. STRINGFELLOW, ADAM G. GINSBURG, JOHN BALLY, & BRIAN A. KEENEY
CASA, Department of Astrophysical and Planetary Sciences, University of Colorado 389-UCB, Boulder, CO 80309, USA
Draft version November 21, 2012

ABSTRACT

We present optical and near-IR spectroscopic observations of the luminous blue variable SN 2009ip during its remarkable photometric evolution of 2012. The spectra sample three key points in the SN 2009ip lightcurve, corresponding to its initial brightening in August (2012-A) and its dramatic rebrightening in early October (2012-B). Based on line fluxes and velocities measured in our spectra, we find a surprisingly low $I(\text{H}\alpha)/I(\text{H}\beta)$ ratio (~ 1.5) in the 2012-B spectra. Such a ratio implies either a rare Case B recombination scenario where $\text{H}\alpha$, but not $\text{H}\beta$, is optically thick, or an extremely high density for the circumstellar material of $n_e > 10^{13} \text{ cm}^{-3}$. The $\text{H}\alpha$ line intensity yields a minimum radiating surface area of $\gtrsim 20,000 \text{ AU}^2$ in $\text{H}\alpha$ at the peak of SN 2009ip's photometric evolution. Combined with the nature of this object's spectral evolution in 2012, a high circumstellar density and large radiating surface area imply the presence of a thin disk geometry around the central star (and, consequently, a possible binary companion), suggesting that the observed 2012-B rebrightening of SN 2009ip can be attributed to the illumination of the disk's inner rim by fast-moving ejecta produced by the underlying events of 2012-A.

Subject headings:

1. INTRODUCTION

The object hereafter referred to as SN 2009ip was discovered on 2009 August 26 (UT dates are used throughout this paper) by Maza et al. (2009) and initially identified as a faint ($\sim 17.9 \pm 0.3$ mag) supernova candidate. Miller et al. (2009) soon noted that historical data revealed the presence of a transient at the SN 2009ip location with evidence of photometric variability stretching back several years, and suggested that this supernova candidate was in fact the outburst of a variable star. Berger et al. (2009) later confirmed this with optical spectroscopy, measuring relatively narrow (FWHM $\sim 550 \text{ km s}^{-1}$) Balmer emission features and a relatively low absolute magnitude of $R \sim -13.7$ and thus identifying SN 2009ip as an outbursting luminous blue variable (LBV) associated with the host galaxy NGC 7259 ($D = 24 \text{ Mpc}$; $z = 0.005714$). Li et al. (2009) reported that this outburst showed a great deal of variability, noting that the LBV had faded to 20.2 mag on 2009 September 11 and then rapidly rebrightened to 18.3 mag on 2009 September 23. Both Smith et al. (2010) and Foley et al. (2011) later analyzed SN 2009ip, identifying it as a very massive ($\gtrsim 60 M_\odot$) star showing evidence of repeated eruptions typical of an LBV based on pre-outburst archival photometry. Smith et al. (2010) also note that the fading and rebrightening behavior seen in the SN 2009ip outburst is extreme but comparable to the behavior of other LBVs.

Spectroscopy of SN 2009ip from Smith et al. (2010) during this first observed eruption event corresponded to a full width at half maximum (FWHM) velocity of $\sim 550 \text{ km s}^{-1}$ in the Balmer emission lines. However, contemporaneous spectroscopy from Foley et al. (2011) measured FWHM velocities of $780\text{--}890 \text{ km s}^{-1}$ in the weeks following the explosion, with later spectra show-

ing broad absorption features with a maximum expansion velocity (measured in the blue wing of the absorption component) of $\sim 4500\text{--}7000 \text{ km s}^{-1}$, larger than the velocities measured for any other LBV eruption except the 1843 eruption of η Carinae (Smith 2008). Spectroscopy from this initial outburst by Pastorello et al. (2012) also indicate a maximum expansion velocity of $5000\text{--}6000 \text{ km s}^{-1}$ for the fast-moving ejected material and FWHM velocities of $700\text{--}800 \text{ km s}^{-1}$, increasing to $1100\text{--}1200 \text{ km s}^{-1}$ at later times as the object faded back to quiescence after rebrightening.

A year after the 2009 outburst, Drake et al. (2010) reported additional activity from SN 2009ip detected in the Siding Spring Survey (SSS) data. An outburst was detected on 2010 July 15, once again followed by a fading period beginning on 2010 September 11 and a second outburst on 2010 September 29, similar to the fading and rebrightening behavior timescale of the 2009 outburst. From a 2010 October 6 spectrum, Pastorello et al. (2012) measure an $\text{H}\alpha$ FWHM velocity of $\sim 1300 \text{ km s}^{-1}$ during this outburst. Drake et al. (2010) also noted that these recent frequent outbursts were similar to the pre-core-collapse outburst of SN 2006jc (e.g. Pastorello et al. 2007, Foley et al. 2007) and stated that such behavior could be indicative of the imminent core-collapse and explosion of SN 2009ip.

Pastorello et al. (2012) present photometric evidence of an additional eruptive phase in mid-2011, extending from May to October and marked by similar luminosity peaks and declines to the 2009 and 2010 outbursts. They obtained two spectra during this period, on 2011 September 2 and September 24. While the first is characterized as a dormant phase of SN 2009ip, with a FWHM velocity of $\sim 940 \text{ km s}^{-1}$ for $\text{H}\alpha$, the second spectrum shows narrower profiles (790 km s^{-1} for $\text{H}\alpha$) and the emergence of strong P Cygni absorption in the Balmer lines. The blue absorption wings of these profiles indicate a maximum

¹ Einstein Fellow; Emily.Levesque@colorado.edu

velocity for the fast-moving material of $\sim 12500 \text{ km s}^{-1}$, the highest outflow velocity ever measured for an LBV eruption.

2012 July 24 marked the discovery of the most recent outburst of SN 2009ip since its 2009 discovery (Drake et al. 2012). Spectroscopy of this new outburst (hereafter referred to as 2012-A) from as early as 2012 August 8 once again show strong broad absorption features, with the blue absorption wings of the Balmer lines extending up to 14000 km s^{-1} and Lorentzian emission profiles with FWHM velocities of $\sim 1380 \text{ km s}^{-1}$ (Pastorello et al. 2012). Additional observations from 2012 August 26 (Vinko et al. 2012), and from 2012 September 16-23 (Smith & Mauerhan 2012a, Mauerhan et al. 2012), also showed Balmer lines with very broad P Cygni absorption profiles corresponding to maximum velocities of $\sim 10000 \text{ km s}^{-1}$ and $\sim 13000 \text{ km s}^{-1}$, respectively. Foley et al. (2012) obtained high-resolution spectroscopy of the outburst from 2012 August 26 and noted prominent Balmer emission features with a somewhat lower FWHM velocity of 640 km s^{-1} , but did not include discussion of any P Cygni absorption components. While these observations were described as characteristic of the spectra and velocities associated with Type IIn supernovae (Smith & Mauerhan 2012a, Mauerhan et al. 2012), contemporaneous photometry initially showed no evidence of any increase in brightness, suggesting luminosities too low to be associated with normal core-collapse SNe (Margutti et al. 2012, Martin et al. 2012). Indeed, photometric monitoring actually showed SN 2009ip fading in mid-September by ~ 1 mag (Martin et al. 2012, Mauerhan et al. 2012, Pastorello et al. 2012, Prieto et al. 2012), similar to the post-outburst fading behavior observed in 2009 and 2010.

However, SN 2009ip began to brighten extremely rapidly ~ 7 days later, on 2012 September 23.60 (Brimacombe 2012). This brightening (hereafter 2012-B) continued over a period of two weeks. It spanned ~ 3 -4 mag, reached an apparent peak or plateau on 2012 October 7, and showed an evolution and maximum luminosity consistent with other Type IIn supernovae (see Prieto et al. 2012 for detailed discussion). During this same time, additional spectroscopy of SN 2009ip showed that the previously-observed P Cygni features had since disappeared (Smith & Mauerhan 2012b, Gall et al. 2012, Bohlson 2012, Pastorello et al. 2012). However, recent spectroscopy from 2012 October 15 and 2012 October 22 suggest a re-emergence of P Cygni profiles with comparable velocities of 11200 - 13100 km s^{-1} (Jha et al. 2012, Childress et al. 2012).

A number of different explanations for the 2012 outburst behavior of SN 2009ip have been presented. Mauerhan et al. (2012) suggest that the 2012-A event likely indicates the initial stages of SN 2009ip undergoing core collapse as a low-luminosity supernova, similar to the faint class of Type II-P's (Pastorello et al. 2007), although they also note that the 2012-A event cannot be completely ruled out as a non-terminal LBV-like eruption, and cite the extreme velocities measured in their 2012 spectra as being implausible for any non-terminal outburst. The 2012-B rebrightening event appears to be the product of the fast SN ejecta catching up to and interacting with slower circumstellar medium (CSM) material ejected by past outbursts; the timescale separat-

ing 2012-A and 2012-B agrees with this model based on measured velocities. The 2012 spectra also resemble other early-time spectra of Type IIn supernovae, with broad-winged Lorentzian profiles produced by electron scattering in the SN-CSM interaction region. The disappearance of the broad-line components is explained by the post-shock formation of an opaque cooling shell that serves to mask the inner high-velocity material. Prieto et al. (2012) share this interpretation, and note that the ~ 1 mag dimming that occurred just before 2012-B is also consistent with the scenario of a fast SN shock breaking out of the SN progenitor's dense CSM material. Furthermore, they note that the apparent low-luminosity nature of the proposed SN should be consistent with a low ^{56}Ni yield, which can potentially be confirmed through long-term monitoring of the SN lightcurve decay.

Alternatively, Pastorello et al. (2012) cite the broad P Cygni profiles and $\sim 12500 \text{ km s}^{-1}$ measured in their 2011 spectra as evidence that LBVs can indeed produce high-velocity blastwaves during non-terminal eruptions. They instead propose that 2012-A is the product of a pulsational pair-instability eruptive event, rather than the core-collapse death of SN 2009ip, stating that the low luminosity of the event is also not consistent with the SN death of such a massive LBV. However, they share the same interpretation of 2012-B as Mauerhan et al. (2012) and Prieto et al. (2012), with the rebrightening and spectroscopic variations being driven by the interaction of the fast-moving 2012-A ejecta material with the star's surrounding dense CSM.

Determining the true nature of SN 2009ip and its 2012 behavior is only possible through continued long-term photometric and spectroscopic monitoring of SN 2009ip and careful in-depth analyses of multi-epoch datasets. Here we present our own optical and near-IR spectroscopic observations of SN 2009ip, obtained at three distinct points in its observed photometric evolution: 2012 August 30 (during the 2012-A outburst), 2012 October 2 (during the 2012-B rapid rebrightening), and 2012 October 9 (after the 2012-B peak/plateau). We detail our acquisition and reduction of the data (§2), and present a series of analyses examining the line profiles and velocities measured in our spectra and the resulting implications for the CSM and effective radiating surface of SN 2009ip during its 2012 evolution (§3). Finally, we consider the potential conclusions that can be drawn from this work and the progress that can be made towards understanding the geometry and underlying nature of this remarkable object (§4).

2. OBSERVATIONS AND REDUCTIONS

We obtained a series of optical and near-infrared spectroscopic observations of SN 2009ip at Apache Point Observatory (APO) during its 2012 period of activity. The dates of our spectroscopic observations are illustrated in Figure 1, along with R- and K-band lightcurves from the literature. Our first observations, in late August, coincide with the 2012-A outburst. Later observations in early October correspond to the 2012-B rebrightening and peak/plateau phases.

2.1. Optical Spectroscopy

Our optical observations were obtained using the medium dispersion ($R \sim 5000$) Dual Imaging Spectro-

graph (DIS) on the APO 3.5m telescope. DIS has separate blue and red cameras with the light split between the two using a dichroic with transition wavelength near 5350Å. The B400 grating was used with the blue camera providing a linear dispersion of $\sim 1.85\text{\AA}/\text{pixel}$. The R300 grating used with the red camera extends from the dichroic cutoff to around $1\text{ }\mu\text{m}$ at $\sim 2.26\text{\AA}/\text{pixel}$ dispersion. All observations were taken with the $1''.5$ slit.

Our first optical spectrum was acquired on 2012 August 30. On this date, SN2009ip was observed during bright Moon near minimum airmass from APO (~ 2.1). We obtained three sequential 1000s exposures under photometric conditions but with variable seeing of $\sim 2'' - 3''$. The flux standard BD+28 4211 (Oke 1990) was observed immediately afterwards at ~ 1.1 airmass. The APO 3.5m telescope is an alt-az design, and has no provision to track the slit at the parallactic angle. Under the marginal conditions experienced on August 30, when the LBV was around ~ 17.5 mag, the slit was left in the E-W orientation and the bright galaxy to the West was used to help facilitate positional guiding during the bright Moon phase. This inevitably resulted in some loss of blue light at the shortest wavelengths.

We next observed SN2009ip on 2012 October 2, after the LBV had undergone an apparent SN-like explosion. Seeing during the DIS exposures (3x240s) observed at an airmass of ~ 2.4 was $\sim 2.3''$. Additional spectra were obtained a week later on 2012 October 9; seeing during these exposures (3x300s) was $\sim 1.''3$ at 2.11 airmass. Both observations were conducted under photometric conditions. In addition, since SN 2009ip was substantially brighter (~ 3 mag) during these observations and guiding was no longer an issue, the slit was rotated to and maintained at 90° to the horizon during the observations, approximating the parallactic angle. Thus, the flux calibration for the blue spectra on these dates should be more accurate than for August 30. Flux standards for the 2012 October 2 and 2012 October 9 observations were BD+25 4655 and BD+33 2642, respectively (Oke 1990).

Extraction and wavelength calibration of the DIS spectra were performed using IRAF tasks, and included bias subtraction, flat fielding using quartz lamps, and wavelength calibration utilizing He, Ne, and Ar lamps that were supplemented by sky lines at the blue end. Flux calibrations for each SN 2009ip observation were performed using standard IDL routines and the observed spectrophotometric standard stars for each corresponding night. When merging the red and blue spectra, preference in duplicated spectral regions was given to the red spectrum because of its higher S/N and overall better flux calibration near the dichroic. No adjustment to the flux of either spectrum was made. The reduced spectra provide wavelength coverage extending from $\sim 3480\text{\AA}$ to the dichroic cutoff ($\sim 5350\text{\AA}$), and from the dichroic cutoff to $\sim 9120\text{\AA}$; these limits are not dictated by the sensitivity of the cameras, but instead by the low observed signal at the blue end and available data for performing the flux calibration using the observed standards at the red end.

2.2. Near-IR Spectroscopy

Near-infrared spectra were obtained on two dates at APO, 2012 August 31 and 2012 October 2. The medium-

resolution cross-dispersed spectrograph TripleSpec (Wilson et al. 2004) was used, providing simultaneous spectral coverage from roughly $0.98\text{--}2.46\text{ }\mu\text{m}$ in 5 spectral orders. This yields a resolving power $R \sim 3200$ using a $1''1 \times 43''$ slit. Exposure times were 180s for 2012 August 31 and 120s for 2012 October 2, and were obtained by nodding along two positions on the slit, facilitating sky subtraction and correction of detector artifacts. Total on-source integration times were 2160s and 600s, respectively.

Sky subtraction, flat fielding, wavelength calibration using OH sky lines, and extraction of the spectra were achieved using a modified version of the software package `xspextool` (originally developed for use with SpeX on the NASA Infrared Telescope Facility; Cushing et al. 2004) redesigned for use with the specific instrument characteristics of TripleSpec. A $2''.5$ aperture extraction was used, with spatial background taken between $3'' - 6''$ with a linear fit. The A0V stars HD 181801 and HD 201202 were used as flux and telluric standards on the respective nights. Telluric corrections and flux calibration were performed using the routine `xtellcor` utilizing the method developed and described by Vacca et al. (2003).

3. ANALYSES

3.1. Identification and Fitting of the Emission Features

Our optical and near-IR spectra are shown in Figure 2. Assuming the continuum can be characterized by a single (dominant) temperature for the emitting region, a blackbody fit can be attempted to our spectra. A 10,000 K blackbody fit for the $>5400\text{\AA}$ regime with the strong lines masked out is shown for each spectrum in Figure 2. This provides a good fit to the 2012 October 9 spectrum above $5400\text{\AA}$, though there is excess emission in the blue. Both the optical and near-infrared spectra from 2012 October 2 show good agreement with a 10,000 K blackbody continuum, although there appears to be some excess emission in the K-band that could indicate the initial stage of dust forming in the circumstellar environment. Further observations are required to confirm the phenomenological explanation for either excess. The early 2012 August 30-31 spectra are not well represented by a 10,000 K blackbody; cooler fits also remain poor. However, the spectra are separated temporally by a day, and no IR photometry has yet been reported for 2012 August 31 which could help with the absolute matching between the optical and IR spectra. The possible loss of blue light due to atmospheric dispersion affects also contributes to uncertainties in the blackbody fitting. Figure 3 shows the flux evolution of our three optical spectra, highlighting the substantial brightening that has occurred in the continuum, while Figure 4 illustrates the similar changes observed in the fluxes of the hydrogen Balmer lines as a function of time. Both the continuum flux and the emission line fluxes increase dramatically from the 2012-A period to the start of the 2012-B rebrightening, and show a further increase in flux between 2012 October 2 and 2012 October 9 as SN 2009ip reached its peak 2012-B brightness.

All three epochs of observations display strong emission lines of *only* the hydrogen Balmer series and helium; key line identifications are listed in Table 1. The integrated emission component of the line flux (F) is given in

Table 1, along with the continuum-subtracted peak line flux (F_{peak}). Fitting of line profiles in our data was done using the `splot` routine in IRAF. These fits were performed for both normalized spectra (to better identify the continuum on either side of the lines) and unnormalized spectra for comparison. The main distinction lies in whether the defined continuum takes into account the uncertain “absorption” components on the blue side of some lines. We have fit the continuum over a longer baseline, focusing on fitting the broad and narrow emission line components only. Line fitting was performed using a single component first, either a Gaussian or a Lorentzian profile. A Lorentzian profile was applied only if it provided a substantial improvement to the fit over that of a Gaussian profile, considering both the narrow and broad (or wing) components. Indeed, a Lorentzian was found to be required in many instances. A Lorentzian profile invariably fits the line profile better when the broad component had a significantly higher velocity compared to the narrow core of the line. When a single Lorentzian fit was inadequate, then a two-component fit comprised of a Gaussian (narrow component) and a Lorentzian (broad component) profile was performed simultaneously. Two-component fits are required for all of the October 9 emission lines (except H δ , which was fit with a Lorentzian), and the October 2 H α , H β , and He I 5876Å lines. For the August 30 spectrum, a single Lorentzian profile adequately fits the H α and H β lines, while a Gaussian profile fits the H γ line. Full-width zero-intensity velocities (v_{FWZI}) attributed to the broad emission component are also listed in Table 1. Evidence for the broad emission component exists for all dates observed, including the 2012 August 30 spectrum coinciding with the 2012-A brightening, and occur in both the hydrogen Balmer and He I lines. The width of the broad component does vary between the epochs observed, apparently increasing during the 2012-B event. The measured FWHM velocities of the narrow emission line components are also listed in Table 1 and typically range from 400-800 km/s at all epochs.

Fitting was also performed for the IR emission lines. Single Gaussian profiles provided adequate fits to the IR lines, which are quite narrow compared to the optical lines. The resulting velocities, both FWZI and FWHM, were substantially smaller than those observed in the optical. The FWHM line widths of only ~ 200 km/s are typical of the winds produced by LBVs.

Figure 5 displays the temporal evolution of the H α , H β , and He I 5876Å emission lines. A broad component of intermediate line widths ranging up to several thousand km s $^{-1}$ is present in the H α profile even during the 2012-A period, but is much less pronounced or even absent in the H β profile for this epoch. The broad component feature becomes progressively more pronounced during the spectral evolution that coincides with the 2012-B re-brightening. This is accompanied by a significant and progressive increase in F_{peak} at each subsequent epoch. The fitted FWHM of the Lorentzian-dominated line profiles does not reflect the substantial increase seen in the broad profile, and remains a measure of the narrow component. In fact, the August 30 FWHM for H α and H β are larger than at subsequent epochs simply because the overall intensity of the line flux is much smaller at that

earlier time. There is a significant increase in the FWZI between the 2012-A and 2012-B events, but little change between the 2012 October 2 and October 9 profiles beyond the substantial rise in F_{peak} . The FWHM is marked in each panel. Perhaps what is most striking in the evolution of the H α and H β profiles is the lack of any substantial Doppler shift between the various epochs, and the consistently symmetric line profiles.

The situation is somewhat different for the evolution of the He I 5876Å line. It is not present in our first epoch of observations during the 2012-A period, but has already become strong before peak light for 2012 October 2, and continues to increase substantially in strength within the following week. This line is asymmetric, depressed on the blue side with a skewed broad red component. It is possible this could be due in part to blending with an unresolved nearby line to the red. However, there is clearly a very broad component by October 9 that extends to a FWZI of ~ 7000 km/s. The degree to which a P Cygni absorption component may contribute to the blue depression is unclear. The progressive increase in the strength of the He I 5876Å line indicates that the temperature at the velocities where this line is forming has increased substantially between each subsequent epoch. This is also evident in the plots of the H α and H β profiles, in which the lines of He I 4922Å, 7065Å, and 7282Å, all of which are weak to nonexistent in the 2012 August 30 spectrum, have continuously increased in strength at later epochs.

3.2. Implications of the Observed Balmer Decrement

The relative line strengths of the H α and H β emission lines provide constraints on the physical conditions in the emitting region. The continuum-subtracted spectrum provides a measure of Balmer line intensities and the Balmer decrement $D = I(\text{H}\alpha)/I(\text{H}\beta)$. In the absence of reddening, Case B recombination (the usual situation in photo-ionized nebulae when Ly α is optically thick but H α is optically thin) results in a ratio of $D \approx 3.0$; D depends weakly on temperature and varies from 3.3 at 2500 K, to 3.05 at 5,000 K, 2.87 at 10,000 K, and 2.76 at 20,000 K (Osterbrock & Ferland 2005). These values apply to HII regions with densities up to $n_e \sim 10^6$ cm $^{-3}$. Reddening can only increase the Balmer decrement since H β is attenuated more than H α by normal interstellar dust, and the reddening to SN 2009ip is low, with $E(B - V) = 0.019$ (Schlegel et al. 1998).

Drake & Ulrich (1980) calculated the expected Balmer decrements for densities between $10^8 < n_e < 10^{15}$ cm $^{-3}$, a range of electron temperatures $5 \times 10^3 < T_e(\text{K}) < 4 \times 10^4$, and a large range in radiation fields and hydrogen line optical depths. Their results show that at densities between $10^8 < n_e < 10^{13}$ cm $^{-3}$, the Balmer decrement increases to values peaking between 10 to 15 around a density $n_e \approx 10^{10}$ to 10^{12} cm $^{-3}$ depending on T_e . The decrement then starts to thermalize due to collisions and decreases to near unity at $n_e \gtrsim 10^{13}$ cm $^{-3}$. A simple estimate shows that the Balmer line intensities should approach a thermalization around this density. For a mostly ionized plasma with a temperature of 10^4 K, the critical density required to thermalize the Balmer lines is roughly $n_{\text{crit}} \sim A_{ul}/\sigma_3 V_e \sim 10^{14}$ cm $^{-3}$ where $A_{ul} = 4.41 \times 10^7$ s $^{-1}$ is the Einstein A coefficient for the $n = 3-2$ transition in hydrogen, $\sigma_3 \sim 6.4 \times 10^{-15}$ is the Bohr cross-

section of hydrogen in its $n = 3$ state, and $V_e \sim 400 \text{ km s}^{-1}$ is the velocity of electrons, the most likely collision partner at 10^4 K .

In SN 2009ip, the $I(H\alpha)/I(H\beta)$ ratio evolved from $D = 3 \pm 0.2$ on 2012 August 30 to $D = 1.4 \pm 0.1$ on 2012 October 2 and 9. Following Drake & Ulrich (1980), such a low Balmer decrement suggests that the line ratios are starting to approach LTE conditions, implying that the bulk of the emission is produced in a very high density plasma with $n_e > 10^{13} \text{ cm}^{-3}$. Alternatively, it is possible that a high optical depth in $H\alpha$ accompanied by a low optical depth in $H\beta$ could permit a relative increase in the brightness of $H\beta$ despite a decrease in the $H\beta$ emissivity (see Drake & Ulrich 1980, Xu et al. 1992).

3.3. Calculating the Radiating Surface Area and Effective Radius

SN 2009ip is located $43''7$ (5.1 kpc in projection) from the nucleus of the Sb spiral NGC 7259 ($z = 0.005714$ which for $H = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$ corresponds to a distance of 24 Mpc; $1'' = 117 \text{ pc}$). Thus, SN 2009ip is located about 5 kpc from the nucleus, far beyond the bright spiral arms of this galaxy. F606W HST images show that the pre-eruption site contains a dim point source, likely the combined light of the LBV and a parent star cluster.

The observed Balmer line fluxes constrain the radiating surface area. The peak $H\alpha$ flux (maximum intensity of the line at its peak, as opposed to the integrated area) from the 2012 October 9 spectrum of SN 2009ip is $I(H\alpha) = 3.05 \times 10^{-14} \text{ ergs s}^{-1} \text{ cm}^{-2} \text{ \AA}^{-1}$, corresponding to a monochromatic $H\alpha$ luminosity of $L(H\alpha) = 2 \times 10^{39} \text{ ergs s}^{-1} \text{ \AA}^{-1}$ at a distance of 24 Mpc. Assuming that this luminosity is produced by a disk of radius $R(H\alpha)$ at a blackbody temperature of $T = 10^4 \text{ K}$, the *minimum* implied radius of the emitting region is given by $R(H\alpha) \gtrsim (L(H\alpha)/4\pi^2 B_\nu(T)\Delta\nu)^{0.5} \gtrsim 1.2 \times 10^{15} \text{ cm}$, or approximately 80 AU. This in turn yields a minimum radiating surface area of $\sim 20000 \text{ AU}^2$. Here, $\Delta\nu = 7 \times 10^{10} \text{ Hz}$ is the bandwidth corresponding to $1\text{\AA}$.

4. CONCLUSIONS AND FUTURE WORK

Recent work on the 2012-A and 2012-B photometric brightenings of SN 2009ip have proposed that the extraordinary brightening of 2012-B can be explained by the interaction of fast-moving ejecta with much slower-moving ejecta from earlier LBV eruptions. The origins of this fast-moving material are unclear, potentially produced either by the core-collapse death of the LBV (Mauerhan et al. 2012, Prieto et al. 2012) or a massive eruption such as a pulsational pair-instability event (Pastorello et al. 2012). Here we focus on the interaction driving the 2012-B rebrightening, and consider two possible geometries:

Shell model: The 2012-B rebrightening traces a rapidly expanding spherical shell powered by the fast-moving ejecta, or

Disk model: The 2012-B rebrightening traces the inner edge of a dense circumstellar excretion disk illuminated by the fast moving ejecta.

In the shell model, earlier LBV mass loss events produce roughly spherical shells expanding at velocities on the order of half the $H\alpha$ FWHM velocity at the time of ejection. If formed during prior 2009 and 2010 erup-

tions, the expected velocities are around 500 to 1000 km s^{-1} and reach radii of $5\text{--}10 \times 10^{15} \text{ cm}$ at the time of the 2012-B rebrightening. Fast ejecta coinciding with the 2012-A brightening ($\sim 13000 \text{ km s}^{-1}$ from Mauerhan et al. 2012), produced by either a core-collapse SN or an exceptionally powerful LBV eruption, would catch up with an older shell at this radius at the precise time scale that we observe, with the 2012-B rebrightening occurring ~ 40 days after 2012-A. If the ejected masses are of order $\sim 1 M_\odot$, the kinetic energy released by the impact of the fast ejecta on the older, slow ejecta could produce the radiated power observed in the 2012-B event.

However, the spectral behavior is inconsistent with this scenario. First, it is hard to understand the change in the Balmer decrement and the large implied density of the $H\alpha$ emitting region during the 2012-B rebrightening. Second, the impact of the fast SN or LBV ejecta is expected to be first seen as a highly blue-shifted component from the near (approaching) side of the slower spherical shell, with a light-crossing time delay of ~ 2 days before showing a corresponding redshifted signature from the rear (receding) side (see, for example, the evolution of the SN 1987A $H\alpha$ spectrum from Kleiser et al. 2011). However, our evolving line profiles in Figure 5 show that the peak of the emission is within 50 km s^{-1} of the rest velocity of NGC 7259, SN 2009ip’s presumed host, consistent with *no* Doppler shift of the strongest emission.

Alternately, in the disk model, the bulk of the emission arises from the radiating inner edge of a circumstellar disk surrounding the LBV. The eruptive behavior of SN 2009ip is quite similar to the observed behavior seen in η Carinae, and long-term photometric and spectroscopic observations of SN 2009ip imply a semi-periodic nature on the order of a year. Combined, this suggests that SN 2009ip could potentially be modeled as a binary system. When the more massive member of a binary evolves off the main sequence and expands, mass transfer onto the companion or Roche lobe overflow can result in a massive circumstellar and circumbinary “excretion” disk. Such disks and rings have been directly observed around many massive stars (e.g. Smith, Bally, & Walawender 2007); examples include SN 1987A, Sher-25 near NGC 3603, and SBW1 (Smith et al. 2012).

In this scenario, the shocks resulting from the interaction of LBV eruptions impact the inner rim of a high-density excretion disk, possibly enhanced by ionizing radiation emitted by the LBV eruption or supernova explosion itself and creating the intense Balmer and helium emission (see Figure 6, top, for a schematic illustration of this geometry). Assuming an inner ring radius of order 10^{15} to 10^{17} cm (ranging from our measured minimum radiating surface radius for SN 2009ip to the typical ring radii seen in Sher-25, SN1987A, and SBW1), when either ionizing radiation or a blastwave hits the inner rim of this very high density disk the resulting emission would be dominated by gas with low radial velocities within 50 km s^{-1} of the host, as observed for SN 2009ip. In the case of the impact of a dense shell from an LBV eruption, or possibly debris from a supernova, entrained material in the fast wind or ejecta flow would produce faint high-velocity emission. Entrained material located along the line of sight to bright continuum sources such as the remnant stellar photosphere, or hot plasma emission from

the inner disk rim, would be seen in absorption, perhaps explaining the high-velocity P-Cygni profiles observed in SN 2009ip (Mauerhan et al. 2012).

Our observations of the SN 2009ip spectrum support this proposed geometry. The peak intensity of the H α line provides a constraint on the minimum surface area radiating this recombination line. Following our results in §3.3, the effective H α radiating surface area on 9 October 2012 in the peak of the H α profile was $A_{rad} \geq 4 \times 10^{30}$ cm² (20,000 AU²). If the disk inner edge has a radius r and thickness h , these parameters have to satisfy the constraint $2\pi rh > A_{rad}$. The observed H α emission could arise from the inner ring edge if the ring has a radius larger than $\sim 10^{15}$ cm, implying a thin ring. The high densities implied by our observed flat Balmer decrement in the spectra taken during the 2012-B rebrightening would therefore be a natural consequence of this circumstellar geometry. The timescale implied by this geometry also coincides perfectly with the delay required for illumination of the disk's inner ring by the 13000 km s⁻¹ 2012-A ejecta. Finally, at such high densities all forbidden line emission would be collisionally suppressed, as observed.

These observations do not necessarily preclude the possibility of a shell model for the circumstellar geometry (see Figure 6, bottom). However, in order for such a geometry to agree with our observations the circumstellar material must fulfill precise conditions that are not commonly observed. The H α optical depth must be greater than 1, while H β must remain optically thin. This situation still describes “Case B” recombination, but yields a scenario where not all H α photons reach the observer, while all H β photons do. At higher optical depth, the H β line becomes trapped and H β photons escape as a pair of Pa α and H α photons in the “Case C” limit, which is not consistent with our observation of low $I(\text{H}\alpha)/I(\text{H}\beta)$.

Within this limb-brightened shell model, it is easier to achieve a large radiating surface area than in the disk model. If we assume the same inner radius as above, $r \sim 300$ AU, the limb-brightened region then only needs to be about 10AU thick in order to provide the observed luminosity.

While this work can constrain the geometry of SN 2009ip's circumstellar environment, the origin of the fast-moving ejecta illuminating the circumstellar material remains unclear. While either geometry is consistent with ejecta produced during the 2012-A event, distinguishing between material produced by a core-collapse SN or an extreme LBV eruption is not possible from current data. Future observations of SN 2009ip's spectral and photometric evolution will reveal whether this geometric model is correct and also ultimately clarify the nature of the 2012-A behavior of SN 2009ip.

We gratefully acknowledge useful discussions and contributions from Charles Danforth, Kevin France, Cynthia Froning, and John Stocke. This work was based on observations obtained with the Apache Point Observatory 3.5-meter telescope, which is owned and operated by the Astrophysical Research Consortium. We thank the support staff at Apache Point for their expertise and assistance during these observations. This research made use of `pyspeckit`, an open-source spectroscopic toolkit for Python hosted at <http://pyspeckit.bitbucket.org>. EML is supported by NASA through Einstein Postdoctoral Fellowship grant number PF0-110075 awarded by the Chandra X-ray Center, which is operated by the Smithsonian Astrophysical Observatory for NASA under contract NAS8-03060. BAK gratefully acknowledges support from NASA grant NNX08AC14G and NSF grant AST1109117.

REFERENCES

- Berger, E., Foley, R., & Ivans, I. 2009, *The Astronomer's Telegram*, 2184, 1
- Bohlsen, T. C. 2012, *The Astronomer's Telegram*, 4477, 1
- Brimacombe, J. 2012, *The Astronomer's Telegram*, 4423, 1
- Childress, R., Scalzo, F., Yuan, B., & Tucker, B. S. 2012, *The Astronomer's Telegram*, 4510, 1
- Cushing, M.C., Vacca, W.D., & Rayner, J.T. 2004, *PASP*, 116, 362
- Drake, S. A. & Ulrich, R. K. 1980, *ApJS*, 42, 351
- Drake, A. J., Howerton, S., McNaught, R., et al. 2012, *The Astronomer's Telegram*, 4334, 1
- Drake, A. J., Prieto, J. L., Djorgovski, S. G., et al. 2010, *The Astronomer's Telegram*, 2897, 1
- Foley, R. J., Smith, N., Ganeshalingam, M., Li, W., Chornock, R., & Filippenko, A. V. 2007, *ApJ*, 657, 105
- Foley, R. J., Berger, E., Fox, Ori, et al. 2011, *ApJ*, 732, 32
- Foley, R. J., Berger, E., Roederer, I. U., & Chomiuk, L. 2012, *The Astronomer's Telegram*, 4338, 1
- Gall, C., Hjorth, J., & Leloudas, G. 2012, *The Astronomer's Telegram*, 4454, 1
- Im, M. 2012, *The Astronomer's Telegram*, 4532, 1
- Jha, S. W., McCully, C., & Patel, B. 2012, *The Astronomer's Telegram*, 4491, 1
- Kleiser, I. K., et al. 2011, *MNRAS*, 415, 372
- Li, W., Smith, N., Miller, A. A., & Filippenko, A. V. 2009, *The Astronomer's Telegram*, 2212, 1
- Margutti, R., Soderberg, A., & Milisavljevic, D. 2012, *The Astronomer's Telegram*, 4414, 1
- Martin, J. C., O'Brien, J., & Hubbell-Thomas, J. 2012, *The Astronomer's Telegram*, 4416, 1
- Mauerhan, J. C., et al. 2012, *MNRAS*, submitted (arXiv:1209.6320)
- Maza, J., Hamuy, M., Antezanas, R., et al. 2009, *Central Bureau Electronic Telegrams*, 1928, 1
- Miller, A. A., Li, W., Nugent, P. E., et al. 2009, *The Astronomer's Telegram*, 2183, 1
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Smith, N., Bally, J., & Walawender, J. 2007, *AJ*, 134, 846
- Smith, N., Miller, A., Li, W., et al. 2010, *AJ*, 139, 1451
- Smith, N. & Mauerhan, J. 2012a, *The Astronomer's Telegram*, 4412, 1
- Smith, N. & Mauerhan, J. 2012b, *The Astronomer's Telegram*, 4427, 1
- Smith, N., Arnett, W. D., Bally, J., Ginsburg, A., & Filipenko, A. V. 2012, *MNRAS*, in press
- Oke, J. B. 1990, *AJ*, 99, 1621
- Osterbrock, D. E. & Ferland, G. J. 2005, *Astrophysics of Gaseous Nebulae and Active Galactic Nuclei* (2nd Edition), University Science Books, 2005
- Pastorello, A., Smartt, S. J., Mattila, S., et al. 2007, *Nature*, 447, 829
- Pastorello, A., Cappellaro, E., Inserra, C., et al. 2012, *ApJ*, submitted (arXiv:1210.3568)
- Prieto, J. L., Brimacombe, J., Drake, A. J., & Howerton, S. 2012, *ApJL*, submitted (arXiv:1210.3347)
- Vacca, W.D., Cushing, M.C., & Rayner, J.T. 2003, *PASP*, 115, 389
- Vinko, J., Wheeler, J. C., Milisavljevic, D., et al. 2012a, *The Astronomer's Telegram*, 4435, 1
- Wilson, J. C., et al. 2004, *SPIE*, 5492, 1295
- Xu, Y., McCray, R., Olivia, E., Randich, S. 1992, *ApJ*, 386, 181




FIG. 1.— R and K photometry of SN 2009ip during late 2012 from Pastorello et al. (2012), Prieto et al. (2012), and Im (2012). The dates of our APO spectroscopy observations are indicated as dashed lines, illustrating their temporal relation to the evolution of the SN 2009ip lightcurve. The earliest detection dates for the 2012-A outburst and 2012-B rebrightening are shown as solid lines. The dotted line shows the average rate of change in the K-band lightcurve from Im (2012)




FIG. 2.— Apache Point Observatory 3.5-m spectrophotometry of SN 2009ip taken with DIS (top) and TripleSpec (bottom), showing data from 2012 August 30–31 (blue), 2012 October 2 (black), and 2012 October 9 (red). The spectra have been corrected to rest-frame wavelengths, and are plotted. The dashed gray lines illustrate the agreement of each spectrum with a 10,000 K blackbody fit.




FIG. 3.— Optical spectrophotometry of SN 2009ip from 2012 August 30–31 (blue), 2012 October 2 (black), and 2012 October 9 (red), showing the temporal evolution of the observed flux.




FIG. 4.— Line profiles of the brightest hydrogen Balmer series emission features, showing changes in these line's fluxes and profiles (on a velocity scale) as a function of time. All spectra have had a linear continuum subtracted. The observations from 2012 August 30 are plotted with both their original fluxes and a multiplicative factor of ten to better illustrate the line profiles. Errors are shown as black bars.




FIG. 5.— Temporal evolution of the $H\alpha$ (top), $H\beta$ (middle), and He I 5876Å (bottom) emission features in our spectra from 2012 August 30–31 (blue), 2012 October 2 (black), and 2012 October 9 (red). The evolution is shown for continuum subtracted line profiles, with the 2012 August 30 data scaled higher by a factor of 5 in order to highlight features at that fainter epoch. The smaller insert shows kinematics over a range of $\pm 3000 \text{ km s}^{-1}$, with the FWHM height & width marked as a dotted line. All lines show an increase in strength with time. Note that in the $H\alpha$ and $H\beta$ panels, the weak He I 4922Å, 7065Å, and 7282Å emission lines also continuously increase in strength at later epochs.




FIG. 6.— Illustration of the two possible circumstellar geometry models that can explain our observations of SN 2009ip; a high-density thin disk model (top) or a limb-brightened shell model where the $H\alpha$ optical depth is greater than 1 while $H\beta$ remains optically thin.

TABLE 1
KEY SN 2009IP EMISSION FEATURES

Species	λ_{rest} (Å)	2012 Aug 30/31 ¹					2012 Oct 2					2012 Oct 9				
		F ¹	F _{peak} ²	v_{FWHM} (km s ⁻¹)	v_{FWZI} (km s ⁻¹)	Fit ³	F	F _{peak}	v_{FWHM} (km s ⁻¹)	v_{FWZI} (km s ⁻¹)	Fit	F	F _{peak}	v_{FWHM} (km s ⁻¹)	v_{FWZI} (km s ⁻¹)	Fit
DIS																
H δ	4101	...	...	...	...	...	3.9	0.26	610	3000	L	6.3	0.72	480	1900	L
H γ	4341	0.30	0.028	730	800	G	7.8	0.61	570	4500	L	15.9	1.33	530	4400	GL
H β	4861	0.80	0.065	750	2000	L	20.1	1.30	540	6000	GL	29.3	2.2	500	6000	GL
He I	5876	...	...	...	...	...	4.5	0.17	870	4800	GL	5.1	0.30	620	3500	GL
H α	6563	4.6	0.216	630	5000	L	33.4	1.78	500	7000	GL	45.7	2.9	450	6000	GL
He I	7065	...	...	...	...	...	0.69	0.070	360	1200	G	2.2	0.19	400	1200	GL
TSpec																
Pa γ	10938	0.84	0.11	200	400	G	0.83	0.10	200	400	G	—	—	—	—	—
Pa β	12818	1.4	0.13	230	600	G	1.6	0.13	230	700	G	—	—	—	—	—
Br γ	21655	0.16	0.014	170	300	G	0.17	0.013	180	300	G	—	—	—	—	—

¹ DIS data from 2012 Aug 30; TripleSpec data from 2012 Aug 31.

² Units of 10⁻¹⁴ ergs s⁻¹ cm⁻² Å⁻¹

³ Fit types; G: Gaussian, L: Lorentzian, GL: Gaussian+Lorentzian